

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
15 June 2000 (15.06.2000)

PCT

(10) International Publication Number
WO 00/34318 A1

(51) International Patent Classification*: C07K 14/435,
C12N 1/00, 1/15, 1/21, 5/10, 15/12, 15/63

117465 (RU). JIANG, Xin; 1133 Rajkovich Way, San Jose,
CA 95120 (DE). DUONG, Tommy; 2491 Glen Elm Way,
San Jose, CA 95148 (US).

(21) International Application Number: PCT/US99/29291

(22) International Filing Date:
10 December 1999 (10.12.1999)

(74) Agent: ADLER, Benjamin, A.; McGregor & Adler, 8011
Candle Ln., Houston, TX 77071 (US).

(25) Filing Language: English

(81) Designated State (national): JP.

(26) Publication Language: English

(84) Designated States (regional): European patent (AT, BE,
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE).

(30) Priority Data:
09/210,330 11 December 1998 (11.12.1998) US
Not furnished 9 December 1999 (09.12.1999) US

Published:
— with international search report

(71) Applicant: CLONTECH LABORATORIES, INC.
[US/US]; 1020 East Meadow Drive, Palo Alto, CA 94303
(US).

(48) Date of publication of this corrected version:
11 April 2002

(72) Inventors: LUKYANOY, Sergey, Anatolievich; ul. Gol-
ubinskaya 13/1-161, Moscow (RU). FRADKOV, Arcady,
Fedorovich; ul. Dnepropetrovskaya, 35/2-14, Moscow,
113570 (RU). LABAS, Yulii Aleksandrovich; ul. Gen-
erala Tyuleneva, 35-416, Moscow, 117465 (RU). MATZ,
Mikhail Vladimirovich; ul. Teplyi stan, 7/2-28, Moscow,

(15) Information about Correction:
see PCT Gazette No. 15/2002 of 11 April 2002, Section II

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: FLUORESCENT PROTEINS FROM NON-BIOLUMINESCENT SPECIES OF CLASS ANTHOZOA, GENES EN-
CODING SUCH PROTEINS AND USES THEREOF

(57) Abstract: The present invention is directed to novel fluorescent proteins from non-bioluminescent organisms from the Class
Anthozoa. Also disclosed are cDNAs encoding the fluorescent proteins.

**FLUORESCENT PROTEINS FROM NON-BIOLUMINESCENT SPECIES
OF CLASS ANTHOZOA, GENES ENCODING SUCH PROTEINS AND
USES THEREOF**

BACKGROUND OF THE INVENTION

Cross-reference to Related Application

This is a divisional application of U.S.S.N. 09/210,330 filed
on December 11, 1998.

Field of the Invention

This invention relates to the field of molecular biology.
More specifically, this invention relates to novel fluorescent proteins,
cDNAs encoding the proteins and uses thereof.

Description of the Related Art

Fluorescence labeling is a particularly useful tool for
marking a protein, cell, or organism of interest. Traditionally, a
protein of interest is purified, then covalently conjugated to a
fluorophore derivative. For *in vivo* studies, the protein-dye complex is
then inserted into cells of interest using micropipetting or a method of
reversible permeabilization. The dye attachment and insertion steps,
however, make the process laborious and difficult to control. An

alternative method of labeling proteins of interest is to concatenate or fuse the gene expressing the protein of interest to a gene expressing a marker, then express the fusion product. Typical markers for this method of protein labeling include β -galactosidase, firefly luciferase and bacterial luciferase. These markers, however, require exogenous substrates or cofactors and are therefore of limited use for *in vivo* studies.

A marker that does not require an exogenous cofactor or substrate is the green fluorescent protein (GFP) of the jellyfish *Aequorea victoria*, a protein with an excitation maximum at 395 nm, a second excitation peak at 475 nm and an emission maximum at 510 nm. GFP is a 238-amino acid protein, with amino acids 65-67 involved in the formation of the chromophore.

Uses of GFP for the study of gene expression and protein localization are discussed in detail by Chalfie et al. in *Science* 263 (1994), 802-805, and Heim et al. in *Proc. Nat. Acad. Sci.* 91 (1994), 12501-12504. Additionally, Rizzuto et al. in *Curr. Biology* 5 (1995), 635-642, discuss the use of wild-type GFP as a tool for visualizing subcellular organelles in cells, while Kaether and Gerdes in *Febs Letters* 369 (1995), 267-271, report the visualization of protein transport along the secretory pathway using wild-type GFP. The expression of GFP in plant cells is discussed by Hu and Cheng in *Febs Letters* 369 (1995), 331-334, while GFP expression in *Drosophila* embryos is described by Davis et al. in *Dev. Biology* 170 (1995), 726-729.

Crystallographic structures of wild-type GFP and the mutant GFP S65T reveal that the GFP tertiary structure resembles a barrel (Ormö et al., *Science* 273 (1996), 1392-1395; Yang, et al., *Nature Biotechnol* 14 (1996), 1246-1251). The barrel consists of beta sheets in a compact structure, where, in the center, an alpha helix containing

the chromophore is shielded by the barrel. The compact structure makes GFP very stable under diverse and/or harsh conditions such as protease treatment, making GFP an extremely useful reporter in general. However, the stability of GFP makes it sub-optimal for determining short-term or repetitive events.

A great deal of research is being performed to improve the properties of GFP and to produce GFP reagents useful and optimized for a variety of research purposes. New versions of GFP have been developed, such as a "humanized" GFP DNA, the protein product of which has increased synthesis in mammalian cells (Haas, et al., *Current Biology* 6 (1996), 315-324; Yang, et al., *Nucleic Acids Research* 24 (1996), 4592-4593). One such humanized protein is "enhanced green fluorescent protein" (EGFP). Other mutations to GFP have resulted in blue-, cyan- and yellow-green light emitting versions. Despite the great utility of GFP, however, other fluorescent proteins with properties similar to or different from GFP would be useful in the art. Novel fluorescent proteins result in possible new colors, or produce pH-dependent fluorescence. Other benefits of novel fluorescent proteins include fluorescence resonance energy transfer (FRET) possibilities based on new spectra and better suitability for larger excitation.

The prior art is deficient in novel fluorescent proteins wherein the DNA coding sequences are known. The present invention fulfills this long-standing need in the art.

SUMMARY OF THE INVENTION

The present invention is directed to DNA sequences encoding fluorescent proteins selected from the group consisting of:

(a) an isolated DNA from an organism from the Class Anthozoa which encodes a fluorescent protein; (b) an isolated DNA which hybridizes to the isolated DNA of (a) and which encodes a fluorescent protein; and (c) an isolated DNA differing from the isolated DNAs of (a) and (b) in codon sequence due to the degeneracy of the genetic code and that encodes a fluorescent protein. Preferably, the DNA is isolated from a non -bioluminescent organism from Class Anthozoa. More preferably, the DNA has the sequence selected from the group consisting of SEQ ID Nos. 55, 57-60, and the fluorescent protein has the amino acid sequence shown in SEQ ID No. 56.

In another embodiment of the present invention, there is provided a vector capable of expressing the DNA of the present invention in a recombinant cell comprising said DNA and regulatory elements necessary for expression of the DNA in the cell. Preferably, the DNA encodes a fluorescent protein having the amino acid sequence shown in SEQ ID No. 56.

In still another embodiment of the present invention, there is provided a host cell transfected with a vector of the present invention, such that the host cell expresses a fluorescent protein. Preferably, the cell is selected from the group consisting of bacterial cells, mammalian cells, plant cells and insect cells.

The present invention is also directed to an isolated and purified fluorescent protein coded for by DNA selected from the group consisting of: (a) isolated DNA from an organism from Class Anthozoa which encodes a fluorescent protein; (b) isolated DNA which hybridizes to the isolated DNA of (a) and which encodes a fluorescent protein; and (c) isolated DNA differing from the isolated DNAs of (a) and (b) in codon sequence due to the degeneracy of the genetic code, and which

encodes a fluorescent protein. Preferably, the protein has the amino acid sequence shown in SEQ ID No. 56.

The present invention is also directed to a DNA sequence encoding a fluorescent protein selected from the group consisting of:

- 5 (a) an isolated DNA which encodes a fluorescent protein, wherein said DNA is from an organism from Class Anthozoa and wherein said organism does not exhibit bioluminescence; (b) an isolated DNA which hybridizes to isolated DNA of (a) and which encodes a fluorescent protein; and (c) an isolated DNA differing from the isolated DNAs of
- 10 (a) and (b) in codon sequence due to degeneracy of the genetic code and which encodes a fluorescent protein. Preferably, the organism is from Sub-class Zoantharia, Order Zoanthidea. More preferably, the organism is from Sub-order Brachycnemina. Even more preferably, the organism is from Family Zoanthidae, Genus Zoanthus. Most
- 15 particularly, the present invention is drawn to a novel fluorescent protein from *Zoanthus sp.*, zFP506.

The present invention is further directed to an amino acid sequence which can be used as a basis for designing an oligonucleotide probe for identification of a DNA encoding a fluorescent protein by

20 means of hybridization, wherein the amino acid sequence is selected from the group consisting of SEQ ID Nos. 3, 5, 8, 11, 12, 14. Preferably, such an oligonucleotide has a nucleotide sequence selected from the group consisting of SEQ ID Nos. 4, 6, 7, 9, 10, 13, 15, 16.

Other and further aspects, features, and advantages of the

25 present invention will be apparent from the following description of the presently preferred embodiments of the invention given for the purpose of disclosure.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the modified strategy of 3'-RACE used to isolate the target fragments. Sequences of the oligonucleotides used are shown in Table 2. Dp1 and Dp2 are the degenerate primers used in the first and second PCR, respectively (see Tables 3 and 4 for the sequences of degenerate primers). In the case of *Zoanthus sp.*, the first degenerate primer used was NGH (SEQ ID No. 4), and the second degenerate primer used was GEGa (SEQ ID No. 6).

Figure 2 shows the excitation and emission spectrum of the novel fluorescent protein from *Zoanthus sp.*, zFP506.

Figure 3 shows functional analysis of destabilized zFP506. It demonstrates that fusion of MODC domain to the zFP506 slightly decreases the fluorescent intensity compared to zFP506 itself and that transient transfection of destabilized zFP506 decreases the fluorescent intensity by 50% after 4-hour treatment with protein synthesis inhibitor cycloheximide (**Figure 3B**) vs. control (**Figure 3A**).

DETAILED DESCRIPTION OF THE INVENTION

As used herein, the term "GFP" refers to the basic green fluorescent protein from *Aequorea victoria*, including prior art versions of GFP engineered to provide greater fluorescence or fluoresce in different colors. The sequence of *Aequorea victoria* GFP (SEQ ID No. 54) has been disclosed in Prasher et al., *Gene* 111 (1992), 229-33.

As used herein, the term "EGFP" refers to mutant variant of GFP having two amino acid substitutions: F64L and S65T (Heim et al., *Nature* 373 (1995), 663-664). The term "humanized" refers to changes

made to the GFP nucleic acid sequence to optimize the codons for expression of the protein in human cells (Yang et al., *Nucleic Acids Research* 24 (1996), 4592-4593).

As used herein, the term "NFP" refers to novel fluorescent protein, and the term "GNFP" refers to green novel fluorescent protein. Specifically, "GNFP" refers to zFP506.

In accordance with the present invention there may be employed conventional molecular biology, microbiology, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. See, e.g., Maniatis, Fritsch & Sambrook, "Molecular Cloning: A Laboratory Manual (1982); "DNA Cloning: A Practical Approach," Volumes I and II (D.N. Glover ed. 1985); "Oligonucleotide Synthesis" (M.J. Gait ed. 1984); "Nucleic Acid Hybridization" (B.D. Hames & S.J. Higgins eds. (1985)); "Transcription and Translation" (B.D. Hames & S.J. Higgins eds. (1984)); "Animal Cell Culture" (R.I. Freshney, ed. (1986)); "Immobilized Cells and Enzymes" (IRL Press, (1986)); B. Perbal, "A Practical Guide To Molecular Cloning" (1984).

A "vector" is a replicon, such as plasmid, phage or cosmid, to which another DNA segment may be attached so as to bring about the replication of the attached segment.

A "DNA molecule" refers to the polymeric form of deoxyribonucleotides (adenine, guanine, thymine, or cytosine) in either single stranded form or a double-stranded helix. This term refers only to the primary and secondary structure of the molecule, and does not limit it to any particular tertiary forms. Thus, this term includes double-stranded DNA found, inter alia, in linear DNA molecules (e.g., restriction fragments), viruses, plasmids, and chromosomes.

A DNA "coding sequence" is a DNA sequence which is transcribed and translated into a polypeptide in vivo when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxyl) terminus. A coding sequence can include, but is not limited to, prokaryotic sequences, cDNA from eukaryotic mRNA, genomic DNA sequences from eukaryotic (e.g., mammalian) DNA, and synthetic DNA sequences. A polyadenylation signal and transcription termination sequence may be located 3' to the coding sequence.

As used herein, the term "hybridization" refers to the process of association of two nucleic acid strands to form an antiparallel duplex stabilized by means of hydrogen bonding between residues of the opposite nucleic acid strands.

The term "oligonucleotide" refers to a short (under 100 bases in length) nucleic acid molecule.

"DNA regulatory sequences", as used herein, are transcriptional and translational control sequences, such as promoters, enhancers, polyadenylation signals, terminators, and the like, that provide for and/or regulate expression of a coding sequence in a host cell.

A "promoter sequence" is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. For purposes of defining the present invention, the promoter sequence is bounded at its 3' terminus by the transcription initiation site and extends upstream (5' direction) to include the minimum number of bases or elements necessary to initiate transcription at levels detectable above background. Within the promoter sequence will be found a

transcription initiation site, as well as protein binding domains responsible for the binding of RNA polymerase. Eukaryotic promoters will often, but not always, contain "TATA" boxes and "CAT" boxes. Various promoters, including inducible promoters, may be used to drive the various vectors of the present invention.

As used herein, the terms "restriction endonucleases" and "restriction enzymes" refer to bacterial enzymes, each of which cut double-stranded DNA at or near a specific nucleotide sequence.

A cell has been "transformed" or "transfected" by exogenous or heterologous DNA when such DNA has been introduced inside the cell. The transforming DNA may or may not be integrated (covalently linked) into the genome of the cell. In prokaryotes, yeast, and mammalian cells for example, the transforming DNA may be maintained on an episomal element such as a plasmid. With respect to eukaryotic cells, a stably transformed cell is one in which the transforming DNA has become integrated into a chromosome so that it is inherited by daughter cells through chromosome replication. This stability is demonstrated by the ability of the eukaryotic cell to establish cell lines or clones comprised of a population of daughter cells containing the transforming DNA. A "clone" is a population of cells derived from a single cell or common ancestor by mitosis. A "cell line" is a clone of a primary cell that is capable of stable growth *in vitro* for many generations.

A "heterologous" region of the DNA construct is an identifiable segment of DNA within a larger DNA molecule that is not found in association with the larger molecule in nature. Thus, when the heterologous region encodes a mammalian gene, the gene will usually be flanked by DNA that does not flank the mammalian genomic DNA in the genome of the source organism. In another example,

heterologous DNA includes coding sequence in a construct where portions of genes from two different sources have been brought together so as to produce a fusion protein product. Allelic variations or naturally-occurring mutational events do not give rise to a
5 heterologous region of DNA as defined herein.

As used herein, the term "reporter gene" refers to a coding sequence attached to heterologous promoter or enhancer elements and whose product may be assayed easily and quantifiably when the construct is introduced into tissues or cells.

10 The amino acids described herein are preferred to be in the "L" isomeric form. The amino acid sequences are given in one-letter code (A: alanine; C: cysteine; D: aspartic acid; E: glutamic acid; F: phenylalanine; G: glycine; H: histidine; I: isoleucine; K: lysine; L: leucine; M: methionine; N: asparagine; P: proline; Q: glutamine; R: arginine; S:
15 serine; T: threonine; V: valine; W: tryptophane; Y: tyrosine; X: any residue). NH₂ refers to the free amino group present at the amino terminus of a polypeptide. COOH refers to the free carboxy group present at the carboxy terminus of a polypeptide. In keeping with standard polypeptide nomenclature, *J Biol. Chem.*, 243 (1969), 3552-
20 59 is used.

The present invention is directed to an isolated DNA selected from the group consisting of: (a) isolated DNA from an organism from the Class Anthozoa which encodes a fluorescent protein; (b) isolated DNA which hybridizes to isolated DNA of (a) and
25 which encodes a fluorescent protein; and (c) isolated DNA differing from the isolated DNAs of (a) and (b) in codon sequence due to the degeneracy of the genetic code, and which encodes a fluorescent protein. Preferably, the DNA has the sequence selected from the group consisting of SEQ ID Nos. 55, 57-60, and the fluorescent protein has the

amino acid sequence shown in SEQ ID No. 56. More preferably, the DNA is non-humanized or humanized zFP506 or N65M.

In another embodiment of the present invention, there is provided a vector capable of expressing the DNA of the present invention in a recombinant cell comprising said DNA and regulatory elements necessary for expression of the DNA in the cell. Specifically, the DNA encodes a fluorescent protein having the amino acid sequence shown in SEQ ID No. 56. Preferably, the vector is constructed by amplifying the DNA and then inserting the amplified DNA into EGFP-N1 backbone, or by fusing different mouse ODC degradation domains such as d1, d2 and d376 to the C-terminal of the DNA and then inserting the fusion DNA into EGFP-N1 backbone.

In still another embodiment of the present invention, there is provided a host cell transfected with the vector of the present invention, which expresses a fluorescent protein of the present invention. Preferably, the cell is selected from the group consisting of bacterial cells, mammalian cells, plant cells, insect cells and yeast cells. A representative example of mammalian cell is HEK 293 cell and an example of bacterial cell is an *E. coli* cell.

The present invention is also directed to a DNA sequence encoding a fluorescent protein selected from the group consisting of: (a) an isolated DNA which encodes a fluorescent protein, wherein said DNA is from an organism from Class Anthozoa and wherein said organism does not exhibit bioluminescence; (b) an isolated DNA which hybridizes to isolated DNA of (a) and which encodes a fluorescent protein; and (c) an isolated DNA differing from the isolated DNAs of (a) and (b) in codon sequence due to degeneracy of the genetic code and which encodes a fluorescent protein. Preferably, the organism is from Sub-class Zoantharia, Order Zoanthidea. More preferably, the

organism is from Sub-order Brachycnemina. Even more preferably, the organism is from Family Zoanthidae, Genus Zoanthus.

The present invention is also directed to an isolated and purified fluorescent protein coded for by DNA selected from the group consisting of: (a) an isolated protein encoded by a DNA which encodes a fluorescent protein wherein said DNA is from an organism from Class Anthozoa and wherein said organism does not exhibit bioluminescence; (b) an isolated protein encoded by a DNA which hybridizes to isolated DNA of (a); and (c) an isolated protein encoded by a DNA differing from the isolated DNAs of (a) and (b) in codon sequence due to degeneracy of the genetic code. Preferably, the isolated and purified fluorescent protein is zFP506.

The present invention is further directed to an amino acid sequence which can be used as a basis for designing an oligonucleotide probe for identification of a DNA encoding a fluorescent protein by means of hybridization, wherein the amino acid sequence is selected from the group consisting of SEQ ID Nos. 3, 5, 8, 11, 12, 14. Preferably, such an oligonucleotide has a nucleotide sequence selected from the group consisting of SEQ ID Nos. 4, 6, 7, 9, 10, 13, 15, 16 and is used as a primer in polymerase chain reaction. Alternatively, it can be used as a probe for hybridization screening of the cloned genomic or cDNA library.

The following examples are given for the purpose of illustrating various embodiments of the invention and are not meant to limit the present invention in any fashion.

EXAMPLE 1

Biological Material

Novel fluorescent proteins were identified from several
5 genera of Anthozoa which do not exhibit any bioluminescence but have
fluorescent color as observed under usual white light or ultraviolet
light. Six species were chosen (see Table 1).

TABLE 1

Anthozoa Species Used in This Study

Species	Area of Origination	Fluorescent Color
Anemonia majano	Western Pacific	bright green tentacle tips
Clavularia sp.	Western Pacific	bright green tentacles and oral disk
Zoanthus sp.	Western Pacific	green-yellow tentacles and oral disk
Discosoma sp. "red"	Western Pacific	orange-red spots oral disk
Discosoma striata	Western Pacific	blue-green stripes on oral disk
Discosoma sp. "magenta"	Western Pacific	faintly purple oral disk
Discosoma sp. "green"	Western Pacific	green spots on oral disk
Anemonia sulcata	Mediterranean	purple tentacle tips

EXAMPLE 2

cDNA Preparation

Total RNA was isolated from the species of interest according to the protocol of Chomczynski and Sacchi (Chomczynski P., et al., *Anal. Biochem.* 162 (1987), 156-159). First-strand cDNA was synthesized starting with 1-3 µg of total RNA using SMART PCR cDNA synthesis kit (CLONTECH) according to the provided protocol with the only alteration being that the "cDNA synthesis primer" provided in the kit was replaced by the primer TN3 (5'- CGCAGTCGACCG(T)₁₃, SEQ ID No. 1) (Table 2). Amplified cDNA samples were then prepared as described in the protocol provided except the two primers used for PCR were the TS primer (5'-AAGCAGTGGTATCAACGCAGAGT, SEQ ID No. 2) (Table 2) and the TN3 primer (Table 2), both in 0.1 µM concentration. Twenty to twenty-five PCR cycles were performed to amplify a cDNA sample. The amplified cDNA was diluted 20-fold in water and 1 µl of this dilution was used in subsequent procedures.

TABLE 2

Oligos Used in cDNA Synthesis and RACE

- 5 TN3: 5'-CGCAGTCGACCG(T)₁₃
(SEQ ID No. 1)
- T7-TN3: 5'-GTAATACGACTCACTATAGGGCCGAGTCGACCG(T)₁₃
(SEQ ID No. 17)
- 10 TS-primer: 5'-AAGCAGTGGTATCAACGCAGAGT
(SEQ ID No. 2)
- T7-TS:
15 5'-GTAATACGACTCACTATAGGGCAAGCAGTGGTATCAACGCAGAGT
(SEQ ID No. 18)
- T7: 5'-GTAATACGACTCACTATAGGGC
(SEQ ID No. 19)
- 20 TS-oligo 5'-AAGCAGTGGTATCAACGCAGAGTACGCrGrGrG
(SEQ ID No. 53)
- 25

EXAMPLE 3

Oligo Design

To isolate fragments of novel fluorescent protein cDNAs,
5 PCR using degenerate primers was performed. Degenerate primers
were designed to match the sequence of the mRNAs in regions that
were predicted to be the most invariant in the family of fluorescent
proteins. Four such stretches were chosen (Table 3) and variants of
degenerate primers were designed. All such primers were directed to
10 the 3'-end of mRNA. All oligos were gel-purified before use. Table 2
shows the oligos used in cDNA synthesis and RACE.

TABLE 3

Key Amino Acid Stretches and Corresponding Degenerate Primers Used for Isolation of Fluorescent Proteins

5

Stretch Position according to A. victoria GFP (7)	Amino Acid Sequence of the Key Stretch	Degenerated Primer Name and Sequence
20-25	GXVNGH (SEQ ID No. 3)	NGH: 5'- GA(C,T) GGC TGC GT(A,T,G,C) AA(T,C) GG(A,T,G) CA (SEQ ID No. 4)
31-35	GEGEG (SEQ ID No. 5) GEGNG (SEQ ID No. 8)	GEGa: 5'- GTT ACA GGT GA(A,G) GG(A,C) GA(A,G) GG (SEQ ID No. 6) GEGb: 5'- GTT ACA GGT GA(A,G) GG(T,G) GA(A,G) GG (SEQ ID No. 7) GNGa: 5'- GTT ACA GGT GA(A,G) GG(A,C) AA(C,T) GG (SEQ ID No. 9) GNGb: 5'- GTT ACA GGT GA(A,G) GG(T,G) AA(C,T) GG (SEQ ID No. 10)
127-131	GMNFP (SEQ ID No. 11) GVNFP (SEQ ID No. 12)	NFP: 5' TTC CA(C,T) GGT (G,A)TG AA(C,T) TT(C,T) CC (SEQ ID NO. 13)
134-137	GPVM (SEQ ID No. 14)	PVMa: 5' CCT GCC (G,A)A(C,T) GGT CC(A,T,G,C) GT(A,C) ATG (SEQ ID NO. 15) PVMb: 5' CCT GCC (G,A)A(C,T) GGT CC(A,T,G,C) GT(G,T) ATG (SEQ ID NO. 16)

EXAMPLE 4

Isolation of 3'-cDNA Fragments of nFPs

The modified strategy of 3'-RACE was used to isolate the target fragments (see Figure 1). The RACE strategy involved two consecutive PCR steps. The first PCR step involved a first degenerate primer (Table 4) and the T7-TN3 primer (SEQ ID No. 17) which has a 3' portion identical to the TN3 primer used for cDNA synthesis (for sequence of T7-TN3, Table 2). The reason for substituting the longer T7-TN3 primer in this PCR step was that background amplification which occurred when using the shorter TN3 primer was suppressed effectively, particularly when the T7-TN3 primer was used at a low concentration (0.1 μ M) (Frohman et al., (1998) *PNAS USA*, 85, 8998-9002). The second PCR step involved the TN3 primer (SEQ ID No. 1, Table 2) and a second degenerate primer (Table 4).

TABLE 4

Combinations of Degenerate Primers for First and Second PCR Resulting in Specific Amplification of 3'-Fragments of nFP cDNA

Species	First Degenerate Primer	Second Degenerate Primer
<i>Anemonia majano</i>	NGH (SEQ ID No. 4)	GNGb (SEQ ID No. 10)
<i>Clavularia</i> sp.	NGH (SEQ ID No. 4)	GEGa (SEQ ID No. 6)
<i>Zoanthus</i> sp.	NGH (SEQ ID No. 4)	GEGa (SEQ ID No. 6)
<i>Discosoma</i> sp. "red"	NGH (SEQ ID No. 4)	GEGa (SEQ ID No. 6), NFP (SEQ ID No. 13) or PVMB (SEQ ID No. 16)
<i>Discosoma striata</i>	NGH (SEQ ID No. 4)	NFP (SEQ ID No. 13)
<i>Anemonia sulcata</i>	NGH (SEQ ID No. 4)	GEGa (SEQ ID No. 6) or NFP (SEQ ID No. 13)

5

The first PCR reaction was performed as follows: 1 μ l of 20-fold
 10 dilution of the amplified cDNA sample was added into the reaction
 mixture containing 1X Advantage KlenTaq Polymerase Mix with
 provided buffer (CLONTECH), 200 μ M dNTPs, 0.3 μ M of first degenerate

primer (Table 4) and 0.1 μ M of T7-TN3 (SEQ ID No. 17) primer in a total volume of 20 μ l. The cycling profile was (Hybaid OmniGene Thermocycler, tube control mode): 1 cycle for 95°C, 10 sec.; 55°C, 1 min.; 72°C, 40 sec; 24 cycles for 95°C, 10 sec.; 62°C, 30 sec.; 72°C, 40 sec. The reaction was then diluted 20-fold in water and 1 μ l of this dilution was added to a second PCR reaction, which contained 1X Advantage KlenTaq Polymerase Mix with the buffer provided by the manufacturer (CLONTECH), 200 μ M dNTPs, 0.3 μ M of the second degenerate primer (Table 4) and 0.1 μ M of TN3 primer. The cycling profile was (Hybaid OmniGene Thermocycler, tube control mode): 1 cycle for 95°C, 10 sec.; 55°C (for GEG/GNG or PVM) or 52°C (for NFP), 1 min.; 72°C, 40 sec; 13 cycles for 95°C, 10sec.; 62°C (for GEG/GNG or PVM) or 58°C (for NFP), 30 sec.; 72°C, 40 sec. The product of PCR was cloned into PCR-Script vector (Stratagene) according to the manufacturer's protocol.

Different combinations of degenerate primers were tried in the first and second PCR reactions on the DNA from each species until a combination of primers was found that resulted in specific amplification--meaning that a pronounced band of expected size (about 650-800 bp for NGH and GEG/GNG and 350-500 bp for NFP and PVM--sometimes accompanied by a few minor bands) was detected on agarose gel after two PCR reactions. The primer combinations of choice for different species of the Class Anthozoa are listed in Table 4. Some other primer combinations also resulted in amplification of fragments of correct size, but the sequence of these fragments showed no homology to the other fluorescent proteins identified or to *Aequorea victoria* GFP.

EXAMPLE 5

Obtaining Full-Length cDNA Copies

Upon sequencing the obtained 3'-fragments of novel
5 fluorescent protein cDNAs, two nested 5'-directed primers were
synthesized for cDNA (Table 5), and the 5' ends of the cDNAs were
then amplified using two consecutive PCRs. In the next PCR reaction,
the novel approach of "step-out PCR" was used to suppress background
amplification. The step-out reaction mixture contained 1x Advantage
10 KlenTaq Polymerase Mix using buffer provided by the manufacturer
(CLONTECH), 200 μ M dNTPs, 0.2 μ M of the first gene-specific primer
(see Table 5), 0.02 μ M of the T7-TS primer (SEQ ID No. 18), 0.1 μ M of
T7 primer (SEQ ID No. 19) and 1 μ l of the 20-fold dilution of the
amplified cDNA sample in a total volume of 20 μ l. The cycling profile
15 was (Hybaid OmniGene Thermocycler, tube control mode): 23-27
cycles for 95°C, 10 sec.; 60°C, 30 sec.; 72°C, 40 sec. The product of
amplification was diluted 50-fold in water and one μ l of this dilution
was added to the second (nested) PCR. The reaction contained 1X
Advantage KlenTaq Polymerase Mix with provided buffer (CLONTECH),
20 200 μ M dNTPs, 0.2 μ M of the second gene-specific primer and 0.1 μ M
of TS primer (SEQ ID No. 2) in a total volume of 20 μ l. The cycling
profile was (Hybaid OmniGene Thermocycler, tube control mode): 12
cycles for 95°C, 10 sec.; 60°C, 30 sec.; 72°C, 40 sec. The product of
amplification was then cloned into pAtlas vector (CLONTECH) according
25 to the manufacturer's protocol.

TABLE 5

Gene-Specific Primers Used for 5'-RACE

Species	First Primer	Second (Nested) Primer
Anemonia majano	5'-GAAATAGTCAGGCATACTGGT (SEQ ID No. 20)	5'-GTCAGGCATAC TGGTAGGAT (SEQ ID No. 21)
Clavularia sp.	5'-CTTGAAATAGTCTGCTATATC (SEQ ID No. 22)	5'-TCTGCTATATC GTCTGGGT (SEQ ID No. 23)
Zoanthus sp.	5'- GTTCTTGAAATAGTCTACTATGT (SEQ ID No. 24)	5'-GTCTACTATGTCTT GAGGAT (SEQ ID No. 25)
Discosoma sp. "red"	5'-CAAGCAAATGGCAAAGGTC (SEQ ID No. 26)	5'-CGGTATTTGTGGCC TTCGTA (SEQ ID No. 27)
Discosoma striata	5'-TTGTCTTCTTCTGCACAAC (SEQ ID No. 28)	5'-CTGCACAACGG GTCCAT (SEQ ID No. 29)
Anemonia sulcata	5'-CCTCTATCTTCATTTCCTGC (SEQ ID No. 30)	5'-TATCTTCATTTCCT GCGTAC (SEQ ID No. 31)
Discosoma sp. "magenta"	5'-TTCAGCACCCCATCACGAG (SEQ ID No. 32)	5'-ACGCTCAGAGCTG GGTTCC (SEQ ID No. 33)
Discosoma sp. "green"	5'-CCCTCAGCAATCCATCACGTTT (SEQ ID No. 34)	5'-ATTATCTCAGTGGA TGGTTC (SEQ ID No. 35)

EXAMPLE 6

Expression of nFP in *E. coli*

5 To prepare a DNA construct for novel fluorescent protein expression, two primers were synthesized for each cDNA: a 5'-directed "downstream" primer with the annealing site located in the 3'-UTR of the cDNA and a 3'-directed "upstream" primer corresponding to the site of translation start site (not including the first ATG codon) (Table 6). Primers with SEQ ID Nos. 41 and 42 were the primers used to prepare the zFP506 DNA. Both primers had 5'-heels coding for a site for a restriction endonuclease; in addition, the upstream primer was designed so as to allow the cloning of the PCR product into the pQE30 vector (Qiagen) in such a way that resulted in the fusion of reading 15 frames of the vector-encoded 6xHis-tag and nFP. The PCR was performed as follows: 1 µl of the 20-fold dilution of the amplified cDNA sample was added to a mixture containing 1x Advantage KlenTaq Polymerase Mix with buffer provided by the manufacturer (CLONTECH), 200 µM dNTPs, 0.2 µM of upstream primer and 0.2 µM of downstream 20 primer, in a final total volume of 20 µl. The cycling profile was (Hybaid OmniGene Thermocycler, tube control mode): 23-27 cycles for 95°C, 10 sec.; 60°C, 30 sec.; 72°C, 40 sec. The product of this amplification step was purified by phenol-chlorophorm extraction and ethanol precipitation and then cloned into pQE30 vector using restriction 25 endonucleases corresponding to the primers' sequence according to standard protocols.

All plasmids were amplified in XL-1 blue *E. coli* and purified by plasmid DNA miniprep kits (CLONTECH). The recombinant clones were selected by colony color, and grown in 3 ml of LB medium

(supplemented with 100 $\mu\text{g/ml}$ of ampicillin) at 37°C overnight. 100 μl of the overnight culture was transferred into 200 ml of fresh LB medium containing 100 $\mu\text{g/ml}$ of ampicillin and grown at 37°C, 200 rpm up to OD_{600} 0.6-0.7. 1 mM IPTG was then added to the culture and
5 incubation was allowed to proceed at 37°C for another 16 hours. The cells were harvested and recombinant protein, which incorporated 6x His tags on the N-terminus, was purified using TALON™ metal-affinity resin according to the manufacturer's protocol (CLONTECH).

TABLE 6

Primers Used to Obtain Full Coding Region of nFPs for Cloning into Expression Construct

Species	Upstream Primer	Downstream Primer
Anemonia majano	5'-acatggatccgctctttcaaca agttatc (SEQ ID No. 36) BamHI	5'-tagtactcgcagcttattcgt tttcagtgaatc (SEQ ID No. 37) XhoI
Clavularia sp.	L: 5'-acatggatcccaacatttttga gaaacg (SEQ ID No. 38) BamHI S: 5'-acatggatccaaagctctaacc accatg (SEQ ID No. 39) BamHI	5'-tagtactcgcagcaacacaa accctcagacaa (SEQ ID No. 40) XhoI
Zoanthus sp.	5'- acatggatccgctcagtcгааг cacggт (SEQ ID No. 41) BamHI	5'-tagtactcgcaggttggaactacat tcctatca (SEQ ID No. 42) XhoI
Discosoma sp. "red"	5'- acatggatccaggtcttccaagaat gttatc (SEQ ID No. 43) BamHI	5'-tagtactcgcaggagccaagttc agcctta (SEQ ID No. 44) XhoI
Discosoma striata	5'- acatggatccagttggtccaagagtgtg (SEQ ID No. 45) BamHI	5'-tagcgcagctctatcatgcctc gtcacct (SEQ ID No. 46) SacI
Anemonia sulcata	5'- acatggatccgcttctcttttaagaagact (SEQ ID No. 47) BamHI	5'-tagtactcgcaggtctctgggagc ggcttg (SEQ ID No. 48) XhoI
Discosoma sp. "magenta"	5'- acatggatccagttgttccaagaatgtgat (SEQ ID No. 49) BamHI	5'-tagtactcgcaggccattacg ctaac (SEQ ID No. 50) XhoI
Discosoma sp. "green"	5'-acatggatccagtgacacttaagaagaatg (SEQ ID No. 51)	5'-tagtactcgcagattcggtttaat gccttg (SEQ ID No. 52)

5

EXAMPLE 7

Novel Fluorescent Proteins and cDNAs Encoding the Proteins

One of the full-length cDNAs encoding fluorescent proteins found is described herein (zFP506). The nucleic acid sequence and deduced amino acid sequence are SEQ ID Nos. 55 and 56, respectively. The spectral properties of zFP506 is listed in Table 7, and the emission and excitation spectra for the ZFP506 is shown in Figure 2.

TABLE 7

Spectral Properties of the Isolated zFP506 nFP

Species:	Zoanthus sp.	Max. Extinction Coefficient:	35,600
nFP Name:	zFP506	Quantum Yield	0.63
Absorbance Max. (nm):	496	Relative Brightness:*	1.02
Emission Max. (nm):	506		

*relative brightness is extinction coefficient multiplied by quantum yield divided by the same value for *A. victoria* GFP.

EXAMPLE 8

Construction of zFP506 Mutant

One mutant of zFP506 was generated, N65M. Compared with wild type zFP506, N65M has the mutation of from "AAC" to "ATG" which results in the corresponding amino acid change from Asparagine

(N) to Methionine (M) at the position of 65. The spectral properties of N65M are listed in Table 8.

TABLE 8

Spectral Properties of the Isolated N65M

	Species:	Zoanthus sp.	Max. Extinction Coefficient:	62,000
10	nFP Name:	N65M	Quantum Yield	0.63
	Absorbance Max. (nm):	496	Relative Brightness:*	1.78
15	Emission Max. (nm):	506		

*relative brightness is extinction coefficient multiplied by quantum yield divided by the same value for *A. victoria* GFP.

EXAMPLE 9

Construction and Functional Analysis of Vectors

Non-humanized zFP506 DNA was amplified via PCR and
25 reconstructed into EGFP-N1 backbone. This vector has the same
multiple cloning sites as EGFP-N1.

Functional test of the generated vector was performed by
transient transfection in 293 cells. 24 hours post transfection,
expression of zFP506 was examined under fluorescent microscope.
30 zFP506 showed good fluorescent intensity and comparable to EGFP-N1.

EXAMPLE 10

Generation of Destabilized zFP506 Vectors as Transcription Reporters

5 Since zFP506 is very stable, it is necessary to generate destabilized versions of zFP506 in order to observe the rapid turnover of the protein. By using the same technology for destabilized EGFP, two destabilized zFP506 vectors were constructed by fusing mouse ODC degradation domain to the C-terminal of zFP506. The d1 version of
10 destabilized zFP506 (SEQ ID No. 57) has three E to A mutations within MODC degradation domain comparing to d2 version (SEQ ID No. 58), therefore result in a shorter half-life of the protein to which MODC degradation domain fused to. Destablized d1zFP506 and d2zFP506 were constructed in EGFP-N1 backbone

EXAMPLE 10

Functional Analysis of Destabilized zFP506

20 Wildtype d1zFP506 was transiently transfected into 293 cells. 24 hours after transfection, CHX was added to stop protein synthesis. After 4 hour treatment, cells were examined under fluorescent microscope. It shows that fusion of MODC domain to the zFP506 slightly decreases the fluorescent intensity compared to zFP506
25 itself. After 4 hour treatment, there is 50% fluorescent intensity decrease (Figure 3B vs. Figure 3A).

EXAMPLE 11

Application of Destabilized d1zFP506 as Transcription Reporters

Destabilized d1zFP506 was constructed into pCRE-d1GNFP
5 and pNF- κ B-d1GNFP vectors. Its expression was regulated under cAMP
response element (CRE) or NF- κ B response element, respectively. These
vectors were transiently transfected into 293 cells, and 24 hours post
transfection, the expression of d1GNFP was induced by Forskolin or
TNF- α . 6 hours after induction, the culture was analysed by FACS. CRE-
10 d1GNFP showed 7 fold of induction in fluorescence intensity, while 4
fold of induction was obtained in NF- κ B-d1GNFP (data not shown). This
demonstrated that the destabilized form of GNFP is applicable as
transcription reporters.

15

EXAMPLE 12

Construction and Functional Test for Humanized zFP506 and Humanized N65M

20 Since mammalian expression is a very popular tool, human
favored codon version is needed for better expression in mammalian
cells. Each piece of human favored codon oligos was linked to form the
full length of wild type and/or mutant zFP506 (hGNFP-zFP506, SEQ ID
No. 59; hGNFP-N65M, SEQ ID No. 60). This humanized zFP506 was
25 constituted into EGFP-N1 backbone.

Any patents or publications mentioned in this specification
are indicative of the levels of those skilled in the art to which the
invention pertains. These patents and publications are incorporated by

reference to the same extent as if each individual publication was specifically and individually indicated to be incorporated by reference.

One skilled in the art will appreciate readily that the present invention is adapted to carry out the objects and obtain the ends and
5 advantages mentioned, as well as those objects and ends inherent therein. The present examples, along with the methods, procedures, treatments, molecules, and specific compounds described herein, are presently representative of preferred embodiments, are exemplary, and are not intended as limitations on the scope of the invention. Changes
10 to the methods and compounds, and other uses, will occur to those skilled in the art and are encompassed within the spirit of the invention as defined by the scope of the claims.

WHAT IS CLAIMED IS:

1. A DNA sequence encoding a fluorescent protein selected from the group consisting of:

5 (a) an isolated DNA which encodes a fluorescent protein, wherein said DNA is from an organism from a Class Anthozoa and wherein said organism does not exhibit bioluminescence;

(b) an isolated DNA which hybridizes to isolated DNA of (a) above and which encodes a fluorescent protein; and

10 (c) an isolated DNA differing from the isolated DNAs of (a) and (b) above in codon sequence due to degeneracy of the genetic code and which encodes a fluorescent protein.

2. The DNA sequence of claim 1, wherein said organism
15 is from Sub-class Zoantharia.

3. The DNA sequence of claim 2, wherein said organism is from Order Zoanthidea.

20 4. The DNA sequence of claim 3, wherein said organism is from Sub-order Brachyncnemina.

5. The DNA sequence of claim 4, wherein said organism is from Family Zoanthidae.

25 6. The DNA sequence of claim 5, wherein said organism is from Genus Zoanthus.

7. A DNA sequence encoding a fluorescent protein selected from the group consisting of:

(a) an isolated DNA which encodes a fluorescent protein having a nucleotide sequence selected from the group consisting of SEQ

5 ID Nos. 55, 57-60;

(b) an isolated DNA which hybridizes to isolated DNA of (a) above and which encodes a fluorescent protein; and

(c) an isolated DNA differing from the isolated DNAs of (a) and (b) above in codon sequence due to degeneracy of the genetic
10 code, and which encodes a fluorescent protein.

8. The DNA sequence of claim 7, wherein said DNA encodes a fluorescent protein having an amino acid sequence shown in
SEQ ID No. 56.

15

9. The DNA sequence of claim 7, wherein said DNA is selected from the group consisting of non-humanized and humanized DNA.

20

10. The DNA sequence of claim 9, wherein said DNA is zFP506 or N65M.

25

11. A vector capable of expressing the DNA sequence of claim 1 in a recombinant cell, wherein said vector comprising said DNA and regulatory elements necessary for expression of the DNA in the cell.

12. The vector of claim 11, wherein said DNA encodes a fluorescent protein having the amino acid sequence shown in SEQ ID No. 56.

5 13. The vector of claim 11, wherein said vector is constructed by amplifying said DNA and then inserting the amplified DNA into EGFP-N1 backbone.

10 14. The vector of claim 13, wherein said DNA is selected from the group consisting of non-humanized and humanized DNA.

15 15. The vector of claim 14, wherein said DNA is zFP506 or N65M.

16. The vector of claim 11, wherein said vector is constructed by fusing different mouse ODC degradation domains to the C-terminal of said DNA and then inserting the fusion DNA into EGFP-N1 backbone.

20 17. The vector of claim 16, wherein said mouse ODC degradation domains are selected from the group consisting of d1, d2 and d376.

25 18. The vector of claim 16, wherein said DNA is selected from the group consisting of non-humanized and humanized DNA.

19. The vector of claim 18, wherein said DNA is zFP506 or N65M.

20. A host cell transfected with the vector of claim 11, wherein said cell is capable of expressing a fluorescent protein.

21. The host cell of claim 20, wherein said cell is selected from the group consisting of bacterial cells, mammalian cells, plant cell, yeast and insect cells.

22. The host cell of claim 21, wherein said mammalian cell is HEK 293 cell.

23. The host cell of claim 21, wherein said bacterial cell is an *E. coli* cell.

24. An isolated and purified fluorescent protein coded for by DNA selected from the group consisting of:

(a) an isolated DNA which encodes a fluorescent protein from an organism from Class Anthozoa, wherein said organism does not exhibit bioluminescence;

(b) an isolated DNA which hybridizes to isolated DNA of (a) above and which encodes a fluorescent protein; and

(c) an isolated DNA differing from the isolated DNAs of (a) and (b) above in codon sequence due to degeneracy of the genetic code and which encodes a fluorescent protein.

25. The isolated and purified fluorescent protein of claim 24, wherein said organism is from Sub-class Zoantharia.

26. The isolated and purified fluorescent protein of claim 25, wherein said organism is from Order Zoanthidea.

27. The isolated and purified fluorescent protein of claim 26, wherein said organism is from Sub-order Brachycnemina.

28. The isolated and purified fluorescent protein of claim 27, wherein said organism is from Family Zoanthidae.

29. The isolated and purified fluorescent protein of claim 28, wherein said organism is from Genus Zoanthus.

30. An isolated and purified fluorescent protein coded for by DNA selected from the group consisting of:

(a) isolated DNA which encodes a fluorescent protein having an amino acid sequence shown in SEQ ID No. 56;

(b) isolated DNA which hybridizes to isolated DNA of (a) above and which encodes a fluorescent protein; and

(c) isolated DNA differing from said isolated DNAs of (a) and (b) above in codon sequence due to degeneracy of the genetic code and which encodes a fluorescent protein.

31. The isolated and purified fluorescent protein of claim 30, wherein said protein is zFP506.

32. An amino acid sequence which can be used as a basis for designing an oligonucleotide probe for identification of a DNA encoding a fluorescent protein by means of hybridization, wherein said sequence is selected from the group consisting of SEQ ID Nos. 3, 5, 8, 11, 12, 14.

33. The amino acid sequence of claim 32, wherein said oligonucleotide has a nucleotide sequence selected from the group consisting of SEQ ID Nos. 4, 6, 7, 9, 10, 13, 15, 16.




FIG. 1
1/3

zFP506




FIG. 2



FIG. 3A



FIG. 3B

3/3

SUBSTITUTE SHEET (RULE 26)

SEQUENCE LISTING

<110> Lukyanov, Sergey A.
 Labas, Yulii A.
 Matz, Mikhail V.
 5 Fradkov, Arcady F.
 Jiang, Xin
 Duong, Tommy
 <120> Fluorescent proteins from non-bioluminescent
 10 species of Class Anthozoa, genes encoding such
 proteins and uses thereof
 <130> D6196D3/PCT
 <141> 1999-12-09
 <150> 09/210,330
 <151> 1998-12-11
 15 <160> 60
 <210> 1
 <211> 25
 <212> DNA
 20 <213> artificial sequence
 <220>
 <221> primer_bind
 <223> primer TN3 used in cDNA synthesis and RACE
 <400> 1
 25 cgccagtcgac cgtttttttt ttttt 25
 <210> 2
 <211> 23
 <212> DNA
 30 <213> artificial sequence
 <220>
 <221> primer_bind
 <223> primer TS used in cDNA synthesis and RACE
 <400> 2
 35 aagcagtggt atcaacgcag agt 23
 <210> 3
 <211> 6
 <212> PRT

<213> *Aequorea victoria*
 <220>
 <222> 21
 <223> amino acid sequence of a key stretch on which
 5 primer NGH is based; Xaa at position 21
 represents unknown
 <400> 3
 Gly Xaa Val Asn Gly His
 5
 10 <210> 4
 <211> 20
 <212> DNA
 <213> artificial sequence
 15 <220>
 <221> primer_bind
 <222> 12
 <223> primer NGH used for isolation of fluorescent
 protein; n at position 12 represents any of the
 20 four bases
 <400> 4
 gayggctgcg tnaayggdca 20
 25 <210> 5
 <211> 5
 <212> PRT
 <213> *Aequorea victoria*
 <220>
 <222> 31...35
 30 <223> amino acid sequence of a key stretch on which
 primers GEGa and GEGb are based
 <400> 5
 Gly Glu Gly Glu Gly
 5
 35 <210> 6
 <211> 20

```

    <212>      DNA
    <213>      artificial sequence
    <220>
    <221>      primer_bind
5    <223>      primer GEGa used for isolation of fluorescent
        protein
    <400>      6

gttacaggtg arggmgargg      20

10    <210>      7
    <211>      20
    <212>      DNA
    <213>      artificial sequence
    <220>
15    <221>      primer_bind
    <223>      primer GEGb used for isolation of fluorescent
        protein
    <400>      7

gttacaggtg arggkgargg      20

20
    <210>      8
    <211>      5
    <212>      PRT
    <213>      Aequorea victoria
25    <220>
    <222>      31...35
    <223>      amino acid sequence of a key stretch on which
        primers GNGa and GNGb are based
    <400>      8

30    Gly Glu Gly Asn Gly
        5

    <210>      9
    <211>      20
35    <212>      DNA
    <213>      artificial sequence
    <220>

```

SEQ 3/23

```

    <221>      primer_bind
    <223>      primer GNGa used for isolation of fluorescent
                protein
    <400>      9
5  gttacaggtg arggmaaygg                                20

    <210>      10
    <211>      20
    <212>      DNA
10  <213>      artificial sequence
    <220>
    <221>      primer_bind
    <223>      primer GNGb used for isolation of fluorescent
                protein
15  <400>      10
    gttacaggtg arggkaaygg                                20

    <210>      11
    <211>      5
20  <212>      PRT
    <213>      Aequorea victoria
    <220>
    <222>      127...131
    <223>      amino acid sequence of a key stretch on which
25  primer NFP is based
    <400>      11
    Gly Met Asn Phe Pro
                5

30  <210>      12
    <211>      5
    <212>      PRT
    <213>      Aequorea victoria
    <220>
35  <222>      127...131
    <223>      amino acid sequence of a key stretch on which
                primer NFP is based

```

SEQ 4/23

	<400>	12	
	Gly Val Asn Phe Pro		
		5	
5	<210>	13	
	<211>	20	
	<212>	DNA	
	<213>	artificial sequence	
	<220>		
10	<221>	primer_bind	
	<223>	primer NFP used for isolation of fluorescent protein	
	<400>	13	
	ttccayggr tgaayttycc		20
15			
	<210>	14	
	<211>	4	
	<212>	PRT	
	<213>	<i>Aequorea victoria</i>	
20	<220>		
	<222>	134...137	
	<223>	amino acid sequence of a key stretch on which primers PVMa and PVMb are based	
	<400>	14	
25	Gly Pro Val Met		
	<210>	15	
	<211>	21	
30	<212>	DNA	
	<213>	artificial sequence	
	<220>		
	<221>	primer_bind	
	<222>	15	
35	<223>	primer PVMa used for isolation of fluorescent protein; n at position 15 represents any of the	

SEQ 5/23

		four bases	
	<400>	15	
	cctgccrayg gtcngtmat g		21
5	<210>	16	
	<211>	21	
	<212>	DNA	
	<213>	artificial sequence	
	<220>		
10	<221>	primer_bind	
	<222>	15	
	<223>	primer PVMb used for isolation of fluorescent protein; n at position 15 represents any of the four bases	
15	<400>	16	
	cctgccrayg gtcngtkat g		21
	<210>	17	
	<211>	47	
20	<212>	DNA	
	<213>	artificial sequence	
	<220>		
	<221>	primer_bind	
	<223>	primer T7-TN3 used in cDNA synthesis and RACE	
25	<400>	17	
	gtaatacgac tcactatagg gccgcagtcg accgtttttt ttttttt		47
	<210>	18	
	<211>	45	
30	<212>	DNA	
	<213>	artificial sequence	
	<220>		
	<221>	primer_bind	
	<223>	primer T7-TS used in cDNA synthesis and RACE	
35	<400>	18	

gtaatacgac tcactatagg gcaagcagtg gtatcaacgc agagt 45

<210> 19

<211> 22

<212> DNA

5 <213> artificial sequence

<220>

<221> primer_bind

<223> primer T7 used in cDNA synthesis and RACE

<400> 19

10 gtaatacgac tcactatagg gc 22

<210> 20

<211> 21

<212> DNA

15 <213> artificial sequence

<220>

<221> primer_bind

<223> gene-specific primer used for 5'-RACE for
Anemonia majano

20 <400> 20

gaaatagtca ggcatactgg t 21

<210> 21

<211> 20

25 <212> DNA

<213> artificial sequence

<220>

<221> primer_bind

<223> gene-specific primer used for 5'-RACE for
Anemonia majano

30 <400> 21

gtcaggcata ctggtaggat 20

<210> 22

35 <211> 21

SEQ 7/23

	<212>	DNA	
	<213>	artificial sequence	
	<220>		
	<221>	primer_bind	
5	<223>	gene-specific primer used for 5'-RACE for <i>Clavularia sp.</i>	
	<400>	22	
	cttgaaatag tctgctatat c		21
10	<210>	23	
	<211>	19	
	<212>	DNA	
	<213>	artificial sequence	
	<220>		
15	<221>	primer_bind	
	<223>	gene-specific primer used for 5'-RACE for <i>Clavularia sp.</i>	
	<400>	23	
	tctgctatat cgtctgggt		19
20	<210>	24	
	<211>	23	
	<212>	DNA	
	<213>	artificial sequence	
	<220>		
25	<221>	primer_bind	
	<223>	gene-specific primer used for 5'-RACE for <i>Zoanthus sp.</i>	
	<400>	24	
30	gttcttgaaa tagtctacta tgt		23
	<210>	25	
	<211>	20	
	<212>	DNA	
35	<213>	artificial sequence	

SEQ 8/23

```

    <220>
    <221>     primer_bind
    <223>     gene-specific primer used for 5'-RACE for
              Zoanthus sp.
5      <400>     25
      gtctactatg tcttgaggat
                                           20

    <210>     26
    <211>     19
10     <212>     DNA
    <213>     artificial sequence
    <220>
    <221>     primer_bind
    <223>     gene-specific primer used for 5'-RACE for
15     <223>     Discosoma sp. "red"
    <400>     26
      caagcaaatg gcaaagggtc
                                           19

    <210>     27
20     <211>     19
    <212>     DNA
    <213>     artificial sequence
    <220>
    <221>     primer_bind
25     <223>     gene-specific primer used for 5'-RACE for
    <223>     Discosoma sp. "red"
    <400>     27
      cggtattgtg gccttcgta
                                           19

30     <210>     28
    <211>     19
    <212>     DNA
    <213>     artificial sequence
    <220>
35     <221>     primer_bind
    <223>     gene-specific primer used for 5'-RACE for

```

SEQ 9/23

Discosoma striata

<400> 28

ttgtcttctt ctgcacaac

19

5 <210> 29
 <211> 17
 <212> DNA
 <213> artificial sequence
 <220>
10 <221> primer_bind
 <223> gene-specific primer used for 5'-RACE for
 Discosoma striata
 <400> 29

ctgcacaacg ggtccat

17

15
 <210> 30
 <211> 20
 <212> DNA
 <213> artificial sequence
20 <220>
 <221> primer_bind
 <223> gene-specific primer used for 5'-RACE for
 Anemonia sulcata
 <400> 30

25 cctctatctt catttcctgc

20

 <210> 31
 <211> 20
 <212> DNA
30 <213> artificial sequence
 <220>
 <221> primer_bind
 <223> gene-specific primer used for 5'-RACE for
 Anemonia sulcata
35 <400> 31

tatcttcatt tcctgcgtac

20

SEQ 10/23

	<210>	32	
	<211>	19	
	<212>	DNA	
5	<213>	artificial sequence	
	<220>		
	<221>	primer_bind	
	<223>	gene-specific primer used for 5'-RACE for <i>Discosoma sp. "magenta"</i>	
10	<400>	32	
	ttcagcaccc	catcacgag	19
	<210>	33	
	<211>	19	
15	<212>	DNA	
	<213>	artificial sequence	
	<220>		
	<221>	primer_bind	
	<223>	gene-specific primer used for 5'-RACE for <i>Discosoma sp. "magenta"</i>	
20	<400>	33	
	acgctcagag	ctgggttcc	19
	<210>	34	
25	<211>	22	
	<212>	DNA	
	<213>	artificial sequence	
	<220>		
	<221>	primer_bind	
30	<223>	gene-specific primer used for 5'-RACE for <i>Discosoma sp. "green"</i>	
	<400>	34	
	ccctcagcaa	tccatcacgt tc	22
35	<210>	35	
	<211>	20	
	<212>	DNA	

SEQ 11/23

	<213>	artificial sequence	
	<220>		
	<221>	primer_bind	
	<223>	gene-specific primer used for 5'-RACE for	
5		<i>Discosoma</i> sp. "green"	
	<400>	35	
	attatctcag tggatgggtc		20
	<210>	36	
10	<211>	31	
	<212>	DNA	
	<213>	artificial sequence	
	<220>		
	<221>	primer_bind	
15	<223>	upstream primer used to obtain full coding region	
		of nFPs from <i>Anemonia majano</i>	
	<400>	36	
	acatggatcc gctctttcaa acaagtttat c		31
20	<210>	37	
	<211>	34	
	<212>	DNA	
	<213>	artificial sequence	
	<220>		
25	<221>	primer_bind	
	<223>	downstream primer used to obtain full coding	
		region of nFPs from <i>Anemonia majano</i>	
	<400>	37	
	tagtactcga gcttattcgt atttcagtga aatc		34
30	<210>	38	
	<211>	29	
	<212>	DNA	
	<213>	artificial sequence	
35	<220>		
	<221>	primer_bind	
	<223>	upstream primer used to obtain full coding region	

SEQ 12/23

of nFPs from *Clavularia sp.*

<400> 38

acatggatcc aacatttttt tgagaaacg 29

5 <210> 39
 <211> 28
 <212> DNA
 <213> artificial sequence
 <220>

10 <221> primer_bind
 <223> upstream primer used to obtain full coding region
 of nFPs from *Clavularia sp.*
 <400> 39

acatggatcc aaagctctaa ccaccatg 28

15 <210> 40
 <211> 31
 <212> DNA
 <213> artificial sequence

20 <220>
 <221> primer_bind
 <223> downstream primer used to obtain full coding
 region of nFPs from *Clavularia sp.*
 <400> 40

25 tagtactcga gcaacacaaa ccctcagaca a 31

<210> 41
 <211> 28
 <212> DNA

30 <213> artificial sequence
 <220>
 <221> primer_bind
 <223> upstream primer used to obtain full coding region
 of nFPs from *Zoanthus sp.*

35 <400> 41

acatggatcc gctcagtaaa agcacggt 28

<210> 42
 <211> 32
 <212> DNA
 5 <213> artificial sequence
 <220>
 <221> primer_bind
 <223> downstream primer used to obtain full coding
 region of nFPs from *Zoanthus sp.*
 10 <400> 42
 tagtactcga ggttggaact acattcttat ca 32
 <210> 43
 <211> 31
 15 <212> DNA
 <213> artificial sequence
 <220>
 <221> primer_bind
 <223> upstream primer used to obtain full coding region
 of nFPs from *Discosoma sp.* "red"
 20 <400> 43
 acatggatcc aggtcttcca agaatgttat c 31
 <210> 44
 25 <211> 29
 <212> DNA
 <213> artificial sequence
 <220>
 <221> primer_bind
 30 <223> downstream primer used to obtain full coding
 region of nFPs from *Discosoma sp.* "red"
 <400> 44
 tagtactcga ggagccaagt tcagcctta 29
 35 <210> 45
 <211> 28
 <212> DNA

SEQ 14/23

<213> artificial sequence
 <220>
 <221> primer_bind
 <223> upstream primer used to obtain full coding region
 5 of nFPs from *Discosoma striata*
 <400> 45
 acatggatcc agttggtcca agagtgtg 28
 <210> 46
 10 <211> 28
 <212> DNA
 <213> artificial sequence
 <220>
 <221> primer_bind
 15 <223> downstream primer used to obtain full coding
 region of nFPs from *Discosoma striata*
 <400> 46
 tagcgagctc tatcatgcct cgtcacct 28
 <210> 47
 20 <211> 31
 <212> DNA
 <213> artificial sequence
 <220>
 25 <221> primer_bind
 <223> upstream primer used to obtain full coding region
 of nFPs from *Anemonia sulcata*
 <400> 47
 acatggatcc gcttcctttt taaagaagac t 31
 30 <210> 48
 <211> 28
 <212> DNA
 <213> artificial sequence
 35 <220>
 <221> primer_bind
 <223> downstream primer used to obtain full coding

SEQ 15/23

region of nFPs from *Anemonia sulcata*

<400> 48

tagtactcga gtccttgagg ggcggcttg 28

5 <210> 49

<211> 30

<212> DNA

<213> artificial sequence

<220>

10 <221> primer_bind

<223> upstream primer used to obtain full coding region
of nFPs from *Discosoma sp. "magenta"*

<400> 49

acatggatcc agttgttcca agaattgat 30

15 <210> 50

<211> 26

<212> DNA

<213> artificial sequence

20 <220>

<221> primer_bind

<223> downstream primer used to obtain full coding
region of nFPs from *Discosoma sp. "magenta"*

<400> 50

25 tagtactcga ggccattacg ctaatc 26

<210> 51

<211> 31

<212> DNA

30 <213> artificial sequence

<220>

<221> primer_bind

<223> upstream primer used to obtain full coding region
of nFPs from *Discosoma sp. "green"*

35 <400> 51

acatggatcc agtgcactta aagaagaaat g 31

SEQ 16/23

<210> 52
 <211> 29
 <212> DNA
 5 <213> artificial sequence
 <220>
 <221> primer_bind
 <223> downstream primer used to obtain full coding
 region of nFPs from *Discosoma* sp. "green"
 10 <400> 52

tagtactcga gattcggttt aatgccttg 29

<210> 53
 <211> 33
 15 <212> DNA
 <213> artificial sequence
 <220>
 <221> primer_bind
 <223> TS-oligo used in cDNA synthesis and RACE
 20 <400> 53

aagcagtggg atcaacgcag agtacgcrgr grg 33

<210> 54
 <211> 238
 25 <212> PRT
 <213> *Aequorea victoria*
 <220>
 <223> amino acid sequence of GFP
 <400> 54

30	Met	Ser	Lys	Gly	Glu	Glu	Leu	Phe	Thr	Gly	Val	Val	Pro	Ile	Leu
					5					10				15	
	Val	Glu	Leu	Asp	Gly	Asp	Val	Asn	Gly	His	Lys	Phe	Ser	Val	Ser
					20					25				30	
	Gly	Glu	Gly	Glu	Gly	Asp	Ala	Thr	Tyr	Gly	Lys	Leu	Thr	Leu	Lys
35					35					40				45	
	Phe	Ile	Cys	Thr	Thr	Gly	Lys	Leu	Pro	Val	Pro	Trp	Pro	Thr	Leu
					50					55				60	

SEQ 17/23

	Val	Thr	Thr	Phe	Ser	Tyr	Gly	Val	Gln	Cys	Phe	Ser	Arg	Tyr	Pro
					65					70					75
	Asp	His	Met	Lys	Gln	His	Asp	Phe	Phe	Lys	Ser	Ala	Met	Pro	Glu
					80					85					90
5	Gly	Tyr	Val	Gln	Glu	Arg	Thr	Ile	Phe	Phe	Lys	Asp	Asp	Gly	Asn
					95					100					105
	Tyr	Lys	Thr	Arg	Ala	Glu	Val	Lys	Phe	Glu	Gly	Asp	Thr	Leu	Val
					110					115					120
	Asn	Arg	Ile	Glu	Leu	Lys	Gly	Ile	Asp	Phe	Lys	Glu	Asp	Gly	Asn
10					125					130					135
	Ile	Leu	Gly	His	Lys	Leu	Glu	Tyr	Asn	Tyr	Asn	Ser	His	Asn	Val
					140					145					150
	Tyr	Ile	Met	Ala	Asp	Lys	Gln	Lys	Asn	Gly	Ile	Lys	Val	Asn	Phe
					155					160					165
15	Lys	Ile	Arg	His	Asn	Ile	Glu	Asp	Gly	Ser	Val	Gln	Leu	Ala	Asp
					170					175					180
	His	Tyr	Gln	Gln	Asn	Thr	Pro	Ile	Gly	Asp	Gly	Pro	Val	Leu	Leu
					185					190					195
	Pro	Asp	Asn	His	Tyr	Leu	Ser	Thr	Gln	Ser	Ala	Leu	Ser	Lys	Asp
20					200					205					210
	Pro	Asn	Glu	Lys	Arg	Asp	His	Met	Val	Leu	Leu	Glu	Phe	Val	Thr
					215					220					225
	Ala	Ala	Gly	Ile	Thr	His	Gly	Met	Asp	Glu	Leu	Tyr	Lys		
					230					235					

25

<210> 55
 <211> 858
 <212> DNA
 <213> *Zoanthus sp.*

30

<220>
 <221> CDS
 <223> cDNA sequence of zFP506
 <400> 55

35

cccaaagcta ctcgactttt gtatcacctt tggggcatca agcgatctgt 50
 tctcaacatg gctcagtc aaacacaggtct aacaaaagaa atgacaatga 100
 aataccgtat ggaaggggtgc gtcgatggac ataaatttgt gatcacggga 150
 gagggcattg gatatccgtt caaagggaaa caggctatta atctgtgtgt 200
 ggtcgaaggt ggaccattgc catttgccga agacatattg tcagctgcct 250

SEQ 18/23

```

ttaactacgg aaacagggtt ttactgaat atcctcaaga catagttgac 300
tatttcaaga actcgtgtcc tgctggatat acatgggaca ggtcttttct 350
ctttgaggat ggagcagttt gcatatgtaa tgcagatata acagtgaatg 400
ttgaagaaaa ctgcatgtat catgagtcca aattttatgg agtgaatttt 450
5  cctgctgatg gacctgtgat gaaaaagatg acagataact gggagccatc 500
ctgcgagaag atcataccag tacctaagca ggggatattg aaaggggatg 550
tctccatgta cctccttctg aaggatggtg ggcgttttacg gtgccaatc 600
gacacagttt acaaagcaaa gtctgtgcc aaaaagatgc cggactggca 650
cttcaccag cataagctca cccgtgaaga ccgcagcgat gctaagaatc 700
10 agaaatggca tctgacagaa catgctattg catccggatc tgcattgccc 750
tgataagaat gtagctccag cattctaata catgtgcttg ccaattatc 800
tgataaatat gtagttgagt tggaaacaga ctagtacaaa taaaagcaca 850
tttaaact 858

```

```

15  <210>      56
    <211>      230
    <212>      PRT
    <213>      Zoanthus sp.
    <220>

```

```

20  <223>      amino acid sequence of zFP506
    <400>      56

```

```

Ala Gln Ser Lys His Gly Leu Thr Lys Glu Met Thr Met Lys Tyr
      5                      10                      15
Arg Met Glu Gly Cys Val Asp Gly His Lys Phe Val Ile Thr Gly
25      20                      25                      30
Glu Gly Ile Gly Tyr Pro Phe Lys Gly Lys Gln Ala Ile Asn Leu
      35                      40                      45
Cys Val Val Glu Gly Gly Pro Leu Pro Phe Ala Glu Asp Ile Leu
      50                      55                      60
30  Ser Ala Ala Phe Asn Tyr Gly Asn Arg Val Phe Thr Glu Tyr Pro
      65                      70                      75
Gln Asp Ile Val Asp Tyr Phe Lys Asn Ser Cys Pro Ala Gly Tyr
      80                      85                      90
Thr Trp Asp Arg Ser Phe Leu Phe Glu Asp Gly Ala Val Cys Ile
35      95                      100                     105
Cys Asn Ala Asp Ile Thr Val Ser Val Glu Glu Asn Cys Met Tyr
      110                     115                     120

```

	His	Glu	Ser	Lys	Phe	Tyr	Gly	Val	Asn	Phe	Pro	Ala	Asp	Gly	Pro
					125					130					135
	Val	Met	Lys	Lys	Met	Thr	Asp	Asn	Trp	Glu	Pro	Ser	Cys	Glu	Lys
					140					145					150
5	Ile	Ile	Pro	Val	Pro	Lys	Gln	Gly	Ile	Leu	Lys	Gly	Asp	Val	Ser
					155					160					165
	Met	Tyr	Leu	Leu	Leu	Lys	Asp	Gly	Gly	Arg	Leu	Arg	Cys	Gln	Phe
					170					175					180
	Asp	Thr	Val	Tyr	Lys	Ala	Lys	Ser	Val	Pro	Arg	Lys	Met	Pro	Asp
10					185					190					195
	Trp	His	Phe	Ile	Gln	His	Lys	Leu	Thr	Arg	Glu	Asp	Arg	Ser	Asp
					200					205					210
	Ala	Lys	Asn	Gln	Lys	Trp	His	Leu	Thr	Glu	His	Ala	Ile	Ala	Ser
					215					220					225
15	Gly	Ser	Ala	Leu	Pro										
					230										

	<210>	57
	<211>	826
20	<212>	DNA
	<213>	artificial sequence
	<220>	
	<222>	497
	<223>	nucleic acid sequence of d1-zFP506, n at position
25		497 represents unknown
	<400>	57

	atggctcag	tcaaagcac	ggttcta	caaaaa	gaaatgaca	atgaaatacc
	tatggaagg	gtcgctcgat	gacataaatt	tgtgatcac	ggagagggc	a
	ttggatatc	gttcaaagg	aaacaggcta	ttaatctgt	gtgtggtcg	aa
30	ggtggaccat	tgccatttgc	cgaagacata	ttgtcagct	gctttaaact	a
	cggaaacagg	gttttctact	aatatcctca	agacatagtt	gactatttca	a
	agaactcgtg	tctgtctgga	tatacatggg	acaggtcttt	tctctttgag	a
	gatggagcag	tttgcata	tgat	ataacagtga	gtgttgaaga	a
	aaactgcatg	tatcatgagt	ccaaatttta	tggagtgaat	tttctctgct	g
35	atggacacgt	gatgaaaaag	atgacagata	actgggagcc	atcctgcgag	a
	aagatcatac	cagtaccta	gcaggggata	ttgaaagggg	atgtctncat	a
	gtacctcctt	ctgaaggatg	gtgggcgttt	acgggtgcaa	ttcgacacag	a

SEQ 20/23

tttacaaagc aaagtctgtg ccaagaaaga tgccgggactg gcacttcac
 cagcataagc tcacccgtga agaccgcagc gatgctaaga atcagaaatg
 gcatctgaca gaacatgcta ttgcatccgg atctgcattg cccaagctta
 gccatggctt cccgccggcg gtggcgggcg aggatgatgg cacgctgccc
 5 atgtcttgtg ccaggagag cgggatggac cgtcaccctg cagcctgtgc
 ttctgctagg atcaatgtgt agatgc 826

<210> 58
 <211> 826
 10 <212> DNA
 <213> artificial sequence
 <220>
 <222> 497
 <223> nucleic acid sequence of d2-zFP506, n at position
 15 497 represents unknown

<400> 58
 atggctcagt caaagcacgg tctaacaaaa gaaatgacaa tgaaataccg 50
 tatggaaggg tgcgtcgatg gacataaatt tgtgatcacg ggagagggca 100
 ttggatatcc gttcaaaagg aaacaggcta ttaatctgtg tgtggtcgaa 150
 20 ggtggaccat tgccatttgc cgaagacata ttgtcagctg cctttaacta 200
 cggaacacagg gttttcactg aatatcctca agacatagtt gactatttca 250
 agaactcgtg tctgtctgga tatacatggg acaggtcttt tctctttgag 300
 gatggagcag tttgcatatg taatgcagat ataacagtga gtgttgaaga 350
 aaactgcagt tatcatgagt ccaaatttta tggagtgaat tttcctgtctg 400
 25 atggacctgt gatgaaaaag atgacagata actgggagcc atcctgcgag 450
 aagatcatac cagtaccta gacggggata ttgaaagggg atgtctncat 500
 gtacctcctt ctgaaggatg gtggcggttt acggtgccaa ttcgacacag 550
 tttacaaagc aaagtctgtg ccaagaaaga tgccgggactg gcacttcac 600
 cagcataagc tcacccgtga agaccgcagc gatgctaaga atcagaaatg 650
 30 gcatctgaca gaacatgcta ttgcatccgg atctgcattg cccaagctta 700
 gccatggctt cccgccggag gtggaggagc aggatgatgg cacgctgccc 750
 atgtcttgtg ccaggagag cgggatggac cgtcaccctg cagcctgtgc 800
 ttctgctagg atcaatgtgt agatgc 826

35 <210> 59
 <211> 696
 <212> DNA

```

    <213>      artificial sequence
    <220>
    <223>      nucleic acid sequence of humanized zFP506
    <400>      59

5  atggcccagt ccaagcacgg cctgaccaag gagatgacca tgaagtaccg 50
   catggagggc tgcgtggacg gccacaagtt cgtgatcacc ggcgagggca 100
   tcggctaccc cttcaagggc aagcaggcca tcaacctgtg cgtggtggag 150
   ggcggccctt tgcccttcgc cgaggacatc ttgtccgccg ccttcaacta 200
   cggaaccgcg gtgttcaccg agtaccacca ggacatcgtc gactacttca 250
10 agaactcctg ccccgccggc tacacctggg accgctcctt cctgttcgag 300
   gacggcgccg tgtgcatctg caacgccgac atcacctgta gcgtggagga 350
   gaactgcatg taccacgagt ccaagttcta cggcgtgaac tccccgccg 400
   acggccccgt gatgaagaag atgaccgaca actgggagcc ctccctgcgag 450
   aagatcatcc ccgtgcccaa gcagggcatc ttgaagggcg acgtgagcat 500
15 gtacctgctg ctgaaggacg gtggccgctt gcgctgccag ttcgacaccg 550
   tgtacaaggc caagtccgtg ccccgcaaga tgcccgaact gcacttcac 600
   cagcacaagc tgacccgcga ggaccgcagc gacgccaaga accagaagtg 650
   gcacctgacc gagcacgcca tcgcctccgg ctccgccttg ccttga 696

20  <210>      60
    <211>      696
    <212>      DNA
    <213>      artificial sequence
    <220>
25  <223>      nucleic acid sequence of humanized N65M
    <400>      60

   atggcccagt ccaagcacgg cctgaccaag gagatgacca tgaagtaccg 50
   catggagggc tgcgtggacg gccacaagtt cgtgatcacc ggcgagggca 100
   tcggctaccc cttcaagggc aagcaggcca tcaacctgtg cgtggtggag 150
30 ggcggccctt tgcccttcgc cgaggacatc ttgtccgccg ccttcatgta 200
   cggaaccgcg gtgttcaccg agtaccacca ggacatcgtc gactacttca 250
   agaactcctg ccccgccggc tacacctggg accgctcctt cctgttcgag 300
   gacggcgccg tgtgcatctg caacgccgac atcacctgta gcgtggagga 350
   gaactgcatg taccacgagt ccaagttcta cggcgtgaac tccccgccg 400
35 acggccccgt gatgaagaag atgaccgaca actgggagcc ctccctgcgag 450
   aagatcatcc ccgtgcccaa gcagggcatc ttgaagggcg acgtgagcat 500
   gtacctgctg ctgaaggacg gcggccgctt gcgctgccag ttcgacaccg 550

```

SEQ 22/23

tgtacaaggc caagtccgtg ccccgcaaga tgcccgactg gcacttcac 600
cagcacaagc tgaccgcgga ggaccgcagc gacgccaaga accagaagtg 650
gcacctgacc gagcacgcca tcgctccgg ctccgccttg ccctga 696

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) Please See Extra Sheet.

US CL Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/320.1, 252.3, 252.33, 325, 410, 254.11, 348, 369, 691; 530/350; 536/23.5

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, P	MATZ et al. Fluorescent proteins from nonbioluminescent Anthozoa species. Nature Biotechnology. October 1999, Volume 17, No. 10, pages 969-973, entire document.	1-33
X, P	DE 197 18 640 A1 (WIEDENMANN) 22 July 1999, entire document.	24-25, 30

☒ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	* I	I
* A* Document defining the general state of the art which is not considered to be of particular relevance	* X*	I
* B* Earlier document published on or after the international filing date	* Y*	I
* L* Document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)		I
* O* Document referring to an oral disclosure, use, exhibition or other means		I
* P* Document published prior to the international filing date but later than the priority date claimed	* A*	I

Date of the actual completion of the international search

23 FEBRUARY 2000

Date of mailing of the international search report

17 MAR 2000

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks

Box PCT

Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

GABRIELE ELISABETH BUGAISKY

Telephone No. (703) 308-0196

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ANDERLUH et al. Cloning, sequencing, and expression of equinatoxin II. Biochemical and Biophysical Research Communications. 1996, Volume 220, No. 2, pages 437-442, entire document.	1-2, 7, 11, 20-21, 23-25, 30
X	MACEK et al. Intrinsic tryptophan fluorescence of equinatoxin II, a pore-forming polypeptide from the sea anemone, Actinia equina L, monitors its interaction with lipid membranes. European Journal of Biochemistry. 1995, Volume 234, pages 329-335, entire document.	24-25, 30
A		1-2, 7, 11, 20-21, 23

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (7):

C07K 14/435; C12N 1/00, 1/15, 1/21, 5/10, 15/12, 15/63

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

435/ 320.1, 252.3, 252.33, 325, 410, 254.11, 348, 369; 530/350; 536/23.5

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

Dialog files 155, 5, 434, 34, 44, 357, 35 (Medline, Biosis, Seisearch, Oceanic Abs., Derwent Biotech Abs, Dissertation Abs.), STN-CAS files Registry, CAPLUS; WEST, files USPT, Derwent WPI
search terms: fluoresce?, bioluminesc?, protein, anthozo?, Zoanth?, Corallimorph?, Discosom?, Coral?, alga, algae,
Caidar?, Invert?, Rhodact?, Actinodisc?, magenta, Clavularia, Zoanthus, Anemonia, majano, anemon?, Zoontha?,
Actinar?, Zoanthid?, Stoloni?, Alcyonar? nqskhgktke/sqp, zfp306, striata, sulcata, brachyconem?, Issaurus

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☒ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.